

**CLINICAL REPORTING FORM**

This form is to be completed **separately** from the client baseline interview. Diagnostic information should be gathered from the respondent's clinical records or diagnostic assessments such as the SCID. The alcohol and drug use rating scale should be completed by a clinician (e.g., client's case manager, psychiatrist, social worker) who is **familiar** with the client's use of alcohol and street drugs.

**DIAGNOSTIC INFORMATION**

**TO BE COMPLETED AT BASELINE INTERVIEW ONLY.**

1. In the client's record, is there evidence of the following?
- 1 Yes      0 No
- |  |   |   |
|--|---|---|
| A. I.Q. below 70   | 1 | 0 |
| B. Prior receipt of services for mental retardation                | 1 | 0 |
| C. Mention of client being of borderline or subnormal intelligence | 1 | 0 |
| D. Head injury   | 1 | 0 |
| E. Spinal cord injury  | 1 | 0 |
| F. Autism  | 1 | 0 |
| G. Cerebral palsy  | 1 | 0 |
| H. Seizure disorder  | 1 | 0 |
| I. Hearing impairment  | 1 | 0 |
| J. Visual impairment   | 1 | 0 |
| K. Specific learning disability                                    | 1 | 0 |
| L. Speech/language impairment                                      | 1 | 0 |
| M. HIV positive  | 1 | 0 |
| N. Attention deficit disorder                                      | 1 | 0 |
| O. Other disability (Specify _____)                                | 1 | 0 |

2. List the respondent's current DSM-IV diagnosis below.

	DSM-IV DIAGNOSIS	DMM-IV CODE
Axis I:	_____	____ . ____
	_____	____ . ____
Axis II:	_____	____ . ____
	_____	____ . ____
Axis III:	_____	____ . ____
	_____	____ . ____
Axis IV:	_____	
	_____	
Axis V:	_____	

Completed by: \_\_\_\_\_

(please print)

ALCOHOL AND DRUG USE

**TO BE COMPLETED AT BASELINE INTERVIEW ONLY.** The alcohol and drug use rating scale should be completed by a clinician (e.g., client's case manager, psychiatrist, social worker) who is **familiar** with the client's use of alcohol and street drugs.

1. How would you rate the respondent's use of alcohol during the past six months?

**Abstinent:** Have not used alcohol during the past 6 months 1

**Use without impairment:** Have used alcohol during the past 6 months, but there is no evidence of persistent or recurrent social, occupational, psychological problems related to use and no evidence of recurrent dangerous use. 2

**Abuse:** Have used alcohol during the past 6 months and there is evidence of persistent or recurrent social, occupational, psychological, or physical problems related to use or evidence of recurrent dangerous use. For example, recurrent alcohol use leads to disruptive behavior and housing problems. Problems have persisted for at least one month. 3

**Dependence:** Have experienced at least three of the following: greater amounts or intervals of use than intended; much of time used obtaining or using alcohol; frequent intoxication or withdrawal interferes with other activities; important activities given up because of alcohol use; continued use despite knowledge of substance-related problems; marked tolerance; characteristic withdrawal symptoms; alcohol taken to relieve or avoid withdrawal symptoms. For example, drinking binges and preoccupation with drinking have caused the respondent to drop out of job training and non-drinking social activities. 4

**Dependence with institutionalization:** Alcohol-related problems are so severe that they make noninstitutional living difficult. For example, constant drinking leads to disruptive behavior and inability to pay rent so that the respondent is frequently reported to the police and seeking hospitalization. 5

2. How would you rate the respondent's use of drugs during the past six months?

**Abstinent:** Have not used drugs during the past 6 months 1

**Use without impairment:** Have used drugs during the past 6 months, but there is no evidence of persistent or recurrent social, occupational, psychological problems related to use and no evidence of recurrent dangerous use. 2

**Abuse:** Have used drugs during the past 6 months and there is evidence of persistent or recurrent social, occupational, psychological, or physical problems related to use or evidence of recurrent dangerous use. For example, recurrent drug use leads to disruptive behavior and housing problems. Problems have persisted for at least one month. 3

**Dependence:** Have experienced at least three of the following: greater amounts or intervals of use than intended; much of time used obtaining or using drugs; frequent intoxication or withdrawal interferes with other activities; important activities given up because of drug use; continued use despite knowledge of substance-related problems; marked tolerance; characteristic withdrawal symptoms; drugs taken to relieve or avoid withdrawal symptoms. For example, binges and preoccupation with drugs have caused the respondent to drop out of job training and non-drug social activities. 4

**Dependence with institutionalization:** Drug-related problems are so severe that they make noninstitutional living difficult. For example, constant drinking leads to disruptive behavior and inability to pay rent so that the respondent is frequently reported to the police and seeking hospitalization. 5

Completed by: \_\_\_\_\_

(please print)